

Cyprinids as a Valuable Source of Essential Fatty Acids for Human Health: A Review¹

W. STEFFENS AND M. WIRTH

*Institute of Freshwater Ecology and Inland Fisheries
Dept. Fish Culture and Fish Pathology
Müggelseedamm 310, D-12561 Berlin-Friedrichshagen
Germany*

Abstract. Cyprinids are important objects of world aquaculture production. Many contain high levels of essential fatty acids of the n-6 and especially the n-3 series which have anti-atherosclerotic efficacy. Phytoplankton is the original source of these fatty acids, and the fatty acid composition of the fish is markedly influenced by the lipid pattern of the food. Freshwater fish such as cyprinids are able to desaturate and elongate greater quantities of dietary C 18 n-6 and n-3 polyunsaturated fatty acids (PUFA) to C 20 and C 22 desaturates. Cyprinids feeding on plankton, e. g., silver carp (*Hypophthalmichthys molitrix*) and bighead carp (*Aristichthys nobilis*), are rich in eicosapentaenoic acid and docosahexaenoic acid. But lipids of grass carp (*Ctenopharyngodon idella*) and common carp (*Cyprinus carpio*) are also characterized by considerable contents of n-3 PUFA. Clinical tests have proven that consumption of cyprinids has beneficial effects on human health, especially for persons with cardiovascular diseases.

Cyprinid culture has increased continuously during the last years, and cyprinids represent 70% of world aquaculture fish production which was 9,417,153 tonnes in 1992 (FAO 1994). Thus consumption of cyprinids is of crucial importance for human nutrition especially in several Asian countries.

The most important cyprinid fish species are silver carp, grass carp, common carp and bighead carp (Table 1) which can be reared successfully in warmwater ponds in polyculture. However, many other cyprinid species are also interesting from the economic point of view.

Significance of essential fatty acids

Two types of essential fatty acids, the n-6 series and the n-3 series, cannot be synthesized by animals or humans and must be supplied in the diet (Horrobin and Manku 1990). Both can be metabolized by the same enzyme sequence (Fig. 1).

Essential fatty acids are required for the normal structures (fluidity, flexibility, permeability) of the membranes and act as precursors of the eicosanoids such as prostaglandins, thromboxanes and leucotrienes. They are involved in cholesterol transport and disposal, and are responsible for

¹Paper presented at the Fourth Asian Fisheries Forum, 16-20 October 1995, Beijing, China.

maintaining the impermeable barrier of the skin, and for regulating permeability in the gut. Since n-6 and n-3 fatty acids interfere with each other's metabolism, neither of them should be consumed in excess.

Whereas the necessity of supply of n-6 fatty acids is well known and undisputed (Horrobin and Manku 1990), knowledge of the significance of n-3 fatty acids became generally accepted in recent years (Lands 1986; Stansby 1990a, 1990b).

Table 1. World aquaculture production (t) of important cyprinid species 1992 (FAO 1994).

Silver carp (<i>Hypophthalmichthys molitrix</i>)	1,616,613
Grass carp (<i>Ctenopharyngodon idella</i>)	1,252,728
Common carp (<i>Cyprinus carpio</i>)	1,022,887
Bighead carp (<i>Aristichthys nobilis</i>)	786,604




Fig 1. Metabolic pathways of the n-6 and n-3 fatty acids.

Today there is no doubt that polyunsaturated n-3 fatty acids have anti-atherosclerotic efficacy (Herold and Kinsella 1986, Pigott and Tucker 1987, Singer 1994, Steffens 1995). This is especially based on

- inhibition of synthesis of the vasoaggressive low density lipoproteins (LDL),
- acceleration of LDL elimination,
- non-influence on vasoprotective high density lipoproteins (HDL) or even enhanced HDL production,
- decrease in total serum triglycerides,
- shifting the eicosanoid balance in favor of the anti-aggregatory fraction,
- reduction of platelet aggregation and prolongation of bleeding time; and
- reduction of blood pressure.

However, several investigations have shown that long-chain n-3 polyunsaturated fatty acids (PUFA) may also have beneficial effects on diseases other than those of the heart and blood vessels, e.g., inflammatory diseases, arthritis, nephritis, Lupus erythematosus, multiple sclerosis, strokes, cancer, skin diseases and asthma.

Fatty Acid Levels in Cyprinid Fish Species

Fish lipids are a significant source of essential fatty acids which have its origin in phytoplankton and reach fish via the food web (Steffens 1989). Lipids of marine fish species are characterized by low levels of linoleic acid (18 : 2 n-6) and linolenic acid (18 : 3 n-3) as well as high levels of long-chain n-3 polyunsaturated fatty acids, especially eicosapentaenoic acid (EPA, 20 : 5 n-3) and docosahexaenoic acid (DHA, 22 : 6 n-3). The ratio of total n-3 to n-6 fatty acids is high, varying between 5 and more than 10.

Freshwater fish, on the other hand, generally contain higher levels of C18 PUFA, but also substantial concentrations of EPA and DHA. In addition the fatty acid composition of freshwater fish is characterized by high proportions of n-6 PUFA, especially linoleic acid and arachidonic acid (AA, 20 : 4 n-6). Therefore the ratio of total n-3 to n-6 fatty acids ranges from 1 to 4.

Unlike marine fish species, freshwater fish are able to desaturate and elongate greater quantities of dietary C18 n-6 and n-3 PUFA to C20 and C22 desaturates (Henderson and Tocher 1987, Sargent et al. 1989).

Cyprinids such as silver carp and bighead carp, which feed on phyto- and zooplankton, are abundant in n-3 PUFA (Table 2). But there are also considerable amounts of arachidonic acid (Steffens et al. 1993).

Muscle lipids of the grass carp, (e.g., an important species for aquaculture which feeds on macrophytes (e.g., filamentous algae, waterweeds and grass), contain high levels of EPA and DHA (Table 3).

The fatty acid composition of the common carp is influenced to a great extent by the fatty acid composition of the feed (Table 4). Compared to supplementary wheat, natural food results in higher levels of essential fatty acids in the muscle triglycerides (Table 5).

Investigations concerning the fatty acid composition of a number of European cyprinids have shown that most of them contain considerable levels of essential fatty acids, especially n-3 PUFA (Table 6 and 7). However, there are differences which may depend on species, nutrition and environmental conditions.

Beneficial Effects of Cyprinid Lipids on Human Health

Experiments to take advantage of the favorable fatty acid composition of cyprinid lipids were carried out in Hungary (Farkas et al. 1987, Joó et al. 1988, Csengeri et al. 1989) and Germany (Steffens et al. 1989, 1991; Wirth et al. 1990b, 1992) with silver carp and bighead carp.

Table 2. Fatty acid composition (%) of the triglycerides of the dorsal and ventral muscle of silver carp and bighead carp (Mieth et al. 1989a, 1989b)

Fatty acids	Silver carp (<i>Hypophthalmichthys molitrix</i>)		Bighead carp (<i>Aristichthys nobilis</i>)	
	Dorsal muscle	Ventral muscle	Dorsal muscle	Ventral muscle
14 : 0	4.5	5.0	2.4	2.8
16 : 0	15.4	15.6	10.8	11.4
16 : 1	10.5	11.2	9.1	9.6
18 : 0	3.2	3.4	2.5	2.6
18 : 1 n - 9	24.8	27.2	27.0	24.5
18 : 2 n - 6	4.3	4.4	3.1	3.5
18 : 3 n - 3	7.0	6.9	7.8	7.8
20 : 1 n - 9	2.5	2.6	2.8	3.7
20 : 4 n - 6	3.3	3.1	3.3	3.1
20 : 5 n - 3	6.6	6.6	10.7	10.8
22 : 1 n - 9	2.9	2.9	3.1	3.1
22 : 5 n - 6	1.4	1.3	1.2	1.2
22 : 5 n - 3	2.0	1.9	2.1	2.0
22 : 6 n - 3	6.0	5.3	9.9	8.9
Σ n - 3	21.6	20.7	30.5	29.5
Σ n - 6	11.0	10.5	9.7	9.3
n - 3 / n - 6	2.0	2.0	3.1	3.2

Table 3. Fatty acid composition (%) of the total lipids of white and red muscle, liver and brain of grass carp (Sýkora and Valenta 1978).

Fatty acids	White muscle	Red muscle	Liver	Brain
16 : 0	18.6	17.4	19.5	18.4
16 : 1	10.9	13.2	8.5	10.1
18 : 0	4.9	3.0	3.9	3.9
18 : 1 n - 9	22.9	26.4	25.4	23.3
18 : 2 n - 6	6.0	6.4	5.1	6.1
18 : 3 n - 3	5.4	9.9	7.2	7.8
20 : 4 n - 6	2.4	3.6	4.3	3.9
20 : 5 n - 3	6.4	3.8	5.0	5.1
22 : 5 n - 3	2.5	2.2	2.6	2.0
22 : 6 n - 3	10.2	6.0	11.6	12.1
Σ n - 3	24.9	23.0	27.2	27.8
Σ n - 6	9.9	11.0	10.6	11.7
n - 3 / n - 6	2.5	2.1	2.6	2.4

Table 4. Fatty acid composition (%) of the muscle triglycerides of common carp (125 - 160 g) fed different diets for 84 d (Steffens et al. 1995).

Fatty acids	Basal diet (control)	Basal diet + 10% corn germ oil	Basal diet + 10% sunflower oil	Basal diet + 10% fish oil	Basal diet + 10% rapeseed oil
14 : 0	1.9	0.9	0.9	4.2	0.9
16 : 0	15.1	10.1	8.1	12.4	7.7
16 : 1	6.2	2.4	2.2	4.9	2.4
18 : 0	1.9	1.1	2.0	1.0	1.0
18 : 1 n - 9	30.0	23.4	24.1	18.9	45.1
18 : 2 n - 6	22.3	51.6	50.9	12.8	25.0
18 : 3 n - 3	5.2	3.1	3.3	8.5	9.8
18 : 4 n - 3	0.5	0.2	tr	3.5	0.2
20 : 4 n - 6	1.3	0.7	1.0	0.2	1.3
20 : 5 n - 3	3.0	1.3	1.3	7.1	1.2
22 : 5 n - 3	1.1	0.4	0.4	1.9	0.5
22 : 6 n - 3	7.0	2.9	2.8	13.7	2.7
Σ n - 3	16.8	7.9	7.8	34.7	14.4
Σ n - 6	25.1	53.9	53.7	14.3	27.4
n - 3 / n - 6	0.7	0.1	0.1	2.4	0.5

Table 5. Fatty acid composition (%) of the muscle triglycerides of market-size carp (> 1,000 g) reared on the basis of natural food only or fed supplementary wheat.

Fatty acids	Carp reared on the basis of natural food	Carp fed supple- mentary wheat
16 : 0	15.9	20.5
16 : 1	11.7	9.8
18 : 1 n - 9	27.8	39.2
18 : 2 n - 6	15.2	10.3
18 : 3 n - 3	8.7	5.8
20 : 4 n - 6	2.2	0.9
20 : 5 n - 3	3.3	1.9
22 : 6 n - 3	1.0	0.4
Σ n - 3	15.3	9.3
Σ n - 6	19.4	12.0
n - 3 / n - 6	0.8	0.8

In a test with spontaneously hypertensive rats, which received a diet containing 10% silver carp oil, systolic blood pressure was lowered and levels of triglycerides and total cholesterol were reduced in blood serum after 8 weeks (Wirth et al. 1990c).

In a comparative experiment, the beneficial effect of silver carp oil containing a higher level of arachidonic acid was more pronounced than the effect of mackerel oil on blood pressure, serum lipids and platelet function (Wirth et al. 1990a, 1992).

In a clinical test, 14 hypertensive patients were put on a 2-week diet of 100 g silver carp meat per day (Wirth et al. 1990b, Steffens et al. 1993). This resulted in a significant drop in systolic blood pressure, decreased levels of triglycerides in the blood plasma, and increased HDL-cholesterol, while phospholipid concentration remained constant (Table 8).

Table 6. Fatty acid composition (%) of the total lipids of the white muscle of several European cyprinid species (Sýkora and Valenta 1978, 1979).

Fatty acids	Rudd (<i>Scardinius erythrophthalmus</i>)	Gudgeon (<i>Gobio gobio</i>)	Dace (<i>Leuciscus leuciscus</i>)	Chub (<i>Leuciscus cephalus</i>)	Bleak (<i>Alburnus alburnus</i>)	Zanthe (<i>Vimba vimba</i>)
16 : 0	18.3	21.2	15.5	20.6	20.2	17.2
16 : 1	4.0	6.2	11.3	2.7	7.9	10.1
18 : 0	5.3	7.5	3.9	4.7	4.4	4.4
18 : 1 n - 9	12.3	15.1	20.4	12.0	28.2	33.2
18 : 2 n - 6	5.3	3.9	5.3	2.7	9.7	6.3
18 : 3 n - 3	5.4	6.2	1.4	4.1	5.1	5.8
20 : 4 n - 6	6.6	8.3	3.0	7.7	5.1	2.1
20 : 5 n - 3	15.8	8.3	9.5	9.3	3.0	4.4
22 : 5 n - 3	3.9	2.2	4.3	3.4	1.9	1.2
22 : 6 n - 3	18.3	13.3	12.0	25.7	6.5	7.1
Σ n - 3	43.9	30.5	28.2	42.9	17.4	19.4
Σ n - 6	12.7	14.0	9.3	11.5	15.9	9.8
n - 3 / n - 6	3.5	2.2	3.0	3.7	1.1	2.0

Table 7. Fatty acid composition (%) of the lipids of several European cyprinid species (Vácha and Tvrzická 1994).

Fatty acids	Tench (<i>Tinca tinca</i>)	Bream (<i>Abramis brama</i>)	Roach (<i>Rutilus rutilus</i>)	Asp (<i>Aspius aspius</i>)	Crucian carp (<i>Carassius carassius</i>)	Barb (<i>Barbus barbus</i>)
16 : 0	17.4	21.6	19.8	15.7	21.2	17.2
16 : 1	16.2	7.3	12.9	12.8	14.6	9.1
18 : 0	2.1	7.2	3.5	3.5	4.5	5.9
18 : 1 n - 9	23.3	32.4	22.7	35.8	27.1	18.3
18 : 2 n - 6	5.2	9.7	8.5	2.4	2.7	2.5
18 : 3 n - 3	10.8	1.3	9.3	3.2	4.8	3.3
20 : 4 n - 6	2.4	3.9	1.2	2.4	1.3	5.1
20 : 5 n - 3	4.7	2.0	3.2	3.6	4.0	6.7
22 : 5 n - 3	1.4	0.7	0.9	1.6	1.2	2.5
22 : 6 n - 3	3.5	3.7	3.2	6.5	4.2	15.8
Σ n - 3	24.5	8.7	19.8	17.6	17.4	31.2
Σ n - 6	8.8	15.1	10.5	5.6	4.7	8.9
n - 3 / n - 6	2.8	0.6	1.9	3.1	3.7	3.5

Table 8. Effect of daily administering of 100 g meat of silver carp for 2- weeks to hypertensive patients on blood pressure (mm Hg) and plasma lipids (mmol·l⁻¹) (Wirth et al. 1990b).

	Pre	Post
Systolic blood pressure	151± 9	136 ± 7*
Diastolic blood pressure	94 ± 7	85 ± 7
Triglycerides	1.82 ± 0.86	1.21 ± 0.53*
HDL-cholesterol	1.52 ± 0.39	1.78 ± 0.44*
Phospholipids	223 ± 35	226 ± 40

* Difference significant (P < 0.05)

Lipids of many cyprinid species exhibit high levels of essential fatty acids, particularly n-3 PUFA. In general there is a balanced ratio of n-3 to n-6 fatty acids. Plankton-feeding species such as silver carp and bighead carp, and also grass carp, and common carp are especially valuable sources of these indispensable fatty acids. Experiments with hypertensive rats and clinical tests have confirmed the beneficial effects of cyprinid fish species in preventing cardiovascular diseases. Thus, cyprinids can be recommended as wholesome food for humans.

References

- Csengeri, I., T. Farkas, I. Joó, F. Müller and J. Oláh. 1989. Eicosapentaenoic acid in freshwater fishes and its role in human nutrition. 40th Ann. Meeting EAAP, Aug. 27-31, 1989, Dublin, Summaries, Vol. I: 287-288.
- FAO. 1994. Aquaculture production 1986 - 1992. FAO Fisheries Circular No. 815, Revision 6.
- Farkas, T., I. Joó, I. Garai, S. Tóth, T. Huszka, J. Oláh, F. Müller, F. Majoros and I. Csengeri. 1987. Az eikozapentaénsav és a dokozahehexaénsav élettani jelentősége és előfordulása hazai halakban. Magyar Táplálkozástudományi Társaság XIV. Vándorgyűlése, Hajdúszoboszló: 77-80.
- Henderson, R.J. and D.R. Tocher. 1987. The lipid composition and biochemistry of freshwater fish. In: Progress in lipid research, Vol. 26 (eds. R.T. Holman, W.W. Christie and H. Sprecher), pp. 281-347. Pergamon Press, New York.
- Herold, P.M. and J.E. Kinsella. 1986. Fish oil consumption and decreased risk of cardiovascular disease: A comparison of findings from animal and human feeding trials. American Journal of Clinical Nutrition 43: 566-598.
- Horrobin, D.F. and M.S. Manku. 1990. Clinical biochemistry of essential fatty acids. In: Omega-6 essential fatty acids. Pathophysiology and roles in clinical medicine (ed. D.F. Horrobin), pp. 21-53. Wiley - Liss, New York.
- Joó, I., T. Farkas, I. Csengeri, I. Garai and S. Tóth. 1988. A new source of eicosapentaenoic acid and its hypolipidaemic effect. 29th International Conference on the Biochemistry of Lipids, 19-22 September 1988, Tokyo. Abstract P61.
- Joó, I., K. Kürti, S. Tóth, T. Farkas and I. Csengeri. 1989. The hypolipidaemic effect of bighead carp diet for long period. FEBS '89, 19th Meeting, 2-7 July 1989, Rome. Abstract TH275.
- Lands, W.E.M. 1986. Fish and human health. Academic Press, Orlando. 170 pp.
- Mieth, G., M. Wirth, M. Friedrich, W. Steffens and U. Lieder. 1989a. Zur Lipidzusammensetzung von Cyprinidenarten. 1. Mitt. Lipidgehalt und Fettsäurespektrum von Silberkarpfen (*Hypophthalmichthys molitrix* [Val.]). Nahrung 33: 91-93.
- Mieth, G., M. Wirth, E. Weigelt, W. Steffens, U. Lieder and M. Friedrich. 1989b. Zur Lipidzusammensetzung von Cyprinidenarten. 2. Mitt. Lipidgehalt und Fettsäurespektrum von Marmorcarpfen (*Aristichthys nobilis*). Nahrung 33: 909-912.
- Pigott, G.M. and B.B. Tucker. 1987. Science opens new horizons for marine lipids in human nutrition. Food Review International 3: 105-138.

- Sargent, J., R.J. Henderson and D.R. Tocher. 1989. The lipids. In: Fish nutrition. 2nd edition (ed. J.E. Halver), pp. 153-228. Academic Press, San Diego.
- Singer, P. 1994. Was sind, wie wirken Omega-3-Fettsäuren? Umschau-Zeitschriftenverlag, Frankfurt am Main. 197 pp.
- Stansby, M.E. 1990a. Nutritional properties of fish oil for human consumption - early developments. In: Fish oils in nutrition (ed. M.E. Stansby), pp. 268-288. Van Nostrand Reinhold, New York.
- Stansby, M.E. 1990b. Nutritional properties of fish oil for human consumption - modern aspects. In: Fish oils in nutrition (ed. M.E. Stansby) pp. 289-308. Van Nostrand Reinhold, New York.
- Steffens, W. 1989. Principles of fish nutrition. Ellis Horwood Ltd., Chichester. 384 pp.
- Steffens, W. 1997. Effects of variation in essential fatty acids in fish feeds on nutritive value of freshwater fish for humans. *Aquaculture* 151: 97-119.
- Steffens, W., U. Lieder, G. Mieth, M. Wirth and M. Friedrich. 1989. Zur Wirkungsweise hochungesättigter Fettsäuren der n-3-Reihe im Lipidstoffwechsel und der Bedeutung phytoplanktonfressender Cypriniden aus Binnengewässern als Eicosapentaensäure-reiche Nahrungsmittel. *Fortschritte der Fischereiwissenschaft* 8: 9-18.
- Steffens, W., U. Lieder, M. Wirth and G. Mieth. 1991. The importance of silver carp and bighead carp as dietetic foodstuffs for prophylaxis and therapy of cardiovascular diseases. *Fischerei-Forschung* 29: 113-114.
- Steffens, W., M. Wirth, G. Mieth and U. Lieder. 1993. Freshwater fish as a source of w-3 polyunsaturated fatty acids and their application to human nutrition. In: Fish nutrition in practice (eds. S.J. Kaushik and P. Luquet), pp. 469-474. INRA, Paris.
- Steffens, W., M. Wirth and B. Rennert. 1995. Effects of adding various oils to the diet on growth, feed conversion and chemical composition of carp (*Cyprinus carpio*). *Archives of Animal Nutrition* 47: 381-389.
- Sýkora, M. and M. Valenta. 1978. Lipidy rybnicích ryb celedi Cyprinidae. *Zivocisná Výroba* 23: 811-824.
- Sýkora, M. and M. Valenta. 1979. Lipidy některých říčních ryb celedi Cyprinidae. *Zivocisná Výroba* 24: 867-880.
- Vácha, F. and E. Tvrzická. 1994. Polynenasycené mastné kyseliny a cholesterol v sladkovodních rybách. *Sb. Ref. Ichtyologické Konf.*, 4.-5. kvetna 1994, VÚRH Vodňany: 43-47.
- Wirth, M., V. Moritz, H. Heine, C. Wagenknecht, I. Berger, G. Mieth, W. Steffens and W. Münckner. 1990a. Vergleichende Untersuchungen zur Wirkung von Makrelen und Silberkarpfenöl auf Blutdruck und Lipidstoffwechsel von spontanhypertensiven Ratten. *Journal of Clinical Chemistry and Clinical Biochemistry* 28: 802-803.
- Wirth, M. C.-U. Wagenknecht, R. Ruzcynski, U. Voigt, W. Gerhard, G. Mieth, W. Steffens and B. Zühl. 1990b. Erste klinische Bewertung der n-3-fettsäurereichen Süßwasserfische Marmor- und Silberkarpfen. *Hochdruck* 10: 116-117.
- Wirth, M., V. Moritz, H. Heine, C. Wagenknecht, G. Mieth, M. Friedrich, W. Steffens and U. Lieder. 1990c. Zur Wirkung von Silberkarpfenöl auf Blutdruck und Lipide von spontanhypertonen Ratten. *Nahrung* 34: 575-578.
- Wirth, M., F. Misselwitz, L. Will-Shahab and G. Mieth. 1992. Effect of eicosapentaenoic acid (EPA) alone and in combination with arachidonic acid (AA) on blood pressure, β -adrenoceptor-stimulated adenylyl cyclase activity, lipids and platelet function. *Symposium on Omega-3 Fatty Acids: Metabolism and Biological Effects*, 15-19 September 1992, Oslo. Abstract 158.